

University of Groningen

Hirschsprung disease - genetics and development

Burzynski, Grzegorz Maciej

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2006

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Burzynski, G. M. (2006). *Hirschsprung disease - genetics and development*. s.n.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

4. References

- Amiel J, Attie T, Jan D, Pelet A, Edery P, Bidaud C, Lacombe D, Tam P, Simeoni J, Flori E, Nihoul-Fekete C, Munnich A, Lyonnet S.
Heterozygous endothelin receptor B (EDNRB) mutations in isolated Hirschsprung disease.
Hum Mol Genet 1996 Mar;5(3):355-7
- Amiel J, Laudier B, Attie-Bitach T, Trang H, de Pontual L, Gener B, Trochet D, Etchevers H, Ray P, Simonneau M, Vekemans M, Munnich A, Gaultier C, Lyonnet S. Polyalanine expansion and frameshift mutations of the paired-like homeobox gene PHOX2B in congenital central hypoventilation syndrome.
Nat Genet 2003;33(4):459-61.
- Amiel J, Lyonnet S
Hirschsprung disease, associated syndromes, and genetics: a review.
J Med Genet 2001 38:729-739
- Anderson RB, Stewart AL, Young HM.
Phenotypes of neural-crest-derived cells in vagal and sacral pathways.
Cell Tissue Res 2006 Jan;323(1):11-25.
- Angrist M, Bolk S, Halushka M, Lapchak PA, Chakravarti A. Germline mutations in glial cell line-derived neurotrophic factor (GDNF) and RET in a Hirschsprung disease patient.
Nat Genet 1996;14:341-4.
- Angrist M, Kauffman E, Slaugenhaupt SA, Matise TC, Puffenberger EG, Washington SS, Lipson A, Cass DT, Reyna T, Weeks DE, et al.
A gene for Hirschsprung disease (megacolon) in the pericentromeric region of human chromosome10.
Nat Genet 1993 Aug;4(4):351-6.
- Attie T, Pelet A, Edery P, Eng C, Mulligan LM, Amiel J, Boutrand L, Beldjord C, Nihoul-Fekete C, Munnich A, et al.
Diversity of RET proto-oncogene mutations in familial and sporadic Hirschsprung disease.
Hum Mol Genet 1995 Aug;4(8):1381-6
- Auricchio A, Casari G, Staiano A, Ballabio A.
Endothelin-B receptor mutations in patients with isolated Hirschsprung disease from a non-inbred population.
Hum Mol Genet 1996 Mar;5(3):351-4.
- Badner J, Sieber W, Garver K, Chakravarti A (1990) A genetic study of Hirschsprung disease.
Am J Hum Genet 46:568-580
- Barembaum M, Bronner-Fraser M.
Early steps in neural crest specification.
Semin Cell Dev Biol 2005 Dec;16(6):642-6

Barlow A, de Graaff E, Pachnis V.

Enteric nervous system progenitors are coordinately controlled by the G protein-coupled receptor EDNRB and the receptor tyrosine kinase RET.

Neuron 2003 Dec 4;40(5):905-16

Batourina E, Tsai S, Lambert S, Sprengle P, Viana R, Dutta S, Hensle T, Wang F, Niederreither K, McMahon AP, Carroll TJ, Mendelsohn CL.

Apoptosis induced by vitamin A signaling is crucial for connecting the ureters to the bladder. Nat Genet 2005 Oct;37(10):1082-9.

Baynash AG, Hosoda K, Giaid A, Richardson JA, Emoto N, Hammer RE, Yanagisawa M.

Interaction of endothelin-3 with endothelin-B receptor is essential for development of epidermal melanocytes and enteric neurons.

Cell 1994 Dec 30;79(7):1277-85.

Bitgood MJ, McMahon AP.

Hedgehog and Bmp genes are coexpressed at many diverse sites of cell-cell interaction in the mouse embryo.

Dev Biol 1995 Nov;172(1):126-38.

Bolk S, Pelet A, Hofstra RM et al. A human model for multigenic inheritance: phenotypic expression in Hirschsprung disease requires both the RET gene and a new 9q31 locus.

Proc Natl Acad Sci U S A 2000;97:268-73.

Borrego S, Wright F, Fernandez R, Williams N, Lopez-Alonso M, Davuluri R, Antinolo G, Eng C A founding locus within the RET proto-oncogene may account for a large proportion of apparently sporadic Hirschsprung disease and a subset of cases of sporadic medullary thyroid carcinoma.

Am J Hum Genet 2003 72:88-100

Brooks AS, Bertoli-Avella AM, Burzynski GM, Breedveld GJ, Osinga J, Boven LG, Hurst JA, Mancini GM, Lequin MH, de Coo RF, Matera I, de Graaff E, Meijers C, Willems PJ, Tibboel D, Oostra BA, Hofstra RM.

Homozygous nonsense mutations in KIAA1279 are associated with malformations of the central and enteric nervous systems.

Am J Hum Genet 2005 Jul;77(1):120-6.

Brooks AS, Oostra BA, Hofstra RM.

Studying the genetics of Hirschsprung's disease: unraveling an oligogenic disorder. Clin Genet 2005 Jan;67(1):6-14

Burns, A. J.

Migration of neural crest-derived enteric nervous system precursor cells to and within the gastrointestinal tract. Int J Dev Biol 2005 49, 143-50.

- Burns, A. J., and Le Douarin, N. M.
Enteric nervous system development: analysis of the selective developmental potentialities of vagal and sacral neural crest cells using quail-chick chimeras.
Anat Rec 2001 262, 16-28.
- Burns AJ, Champeval D, Le Douarin NM.
Sacral neural crest cells colonise aganglionic hindgut in vivo but fail to compensate for lack of enteric ganglia.
Dev Biol 2000 Mar 1;219(1):30-43
- Burns AJ, Douarin NM.
The sacral neural crest contributes neurons and glia to the post-umbilical gut: spatiotemporal analysis of the development of the enteric nervous system.
Development 1998 Nov;125(21):4335-47
- Burzynski GM, Nolte IM, Osinga J, Ceccherini I, Twigt B, Maas S, Brooks A, Verheij J, Plaza Menacho I, Buys CHCM, Hofstra RMW
Localizing a putative mutation as the major contributor to the development of sporadic Hirschsprung disease to the RET genomic sequence between the promoter region and exon 2. *Eur J Hum Genet* 2004 12:604-612
- Cacheux V, Dastot-Le Moal F, Kaariainen H, Bondurand N, Rintala R, Boissier B, Wilson M, Mowat D, Goossens M.
Loss-of-function mutations in SIP1 Smad interacting protein 1 result in a syndromic Hirschsprung disease.
Hum Mol Genet 2001 Jul 1;10(14):1503-10.
- Carrasquillo M, McCallion A, Puffenberg E, Kashuk C, Nouri N, Chakravarti A.
Genome-wide association study and mouse model identify interaction between RET and EDNRB pathways in Hirschsprung disease.
Nat Genet 2002;32:237-244.
- Chakravarti, A and Lyonnet S.
Hirschsprung's disease. In *The Metabolic and Molecular Bases of Inherited Diseases* (ed. C. R. Scriver, A. L. Beaudet, D. Valle, W. S. Sly, B. Childs, K. Kinzler and B. Vogelstein), 2001 pp. 6231-6255. New York: McGraw-Hill.
- Chalazonitis A, D'Autreaux F, Guha U, Pham TD, Faure C, Chen JJ, Roman D, Kan L, Rothman TP, Kessler JA, Gershon MD.
Bone morphogenetic protein-2 and -4 limit the number of enteric neurons but promote development of a TrkC-expressing neurotrophin-3-dependent subset.
J Neurosci 2004 Apr 28;24(17):4266-82.
- Cornell RA, Eisen JS.
Notch in the pathway: the roles of Notch signaling in neural crest development.
Semin Cell Dev Biol 2005 Dec;16(6):663-72

Creuzet S, Couly G, Le Douarin NM.

Patterning the neural crest derivatives during development of the vertebrate head: insights from avian studies.

J Anat 2005 Nov;207(5):447-59

Davidson B, Goldberg I, Tell L, Vigdorchik S, Baekelandt M, Berner A, Kristensen GB, Reich R, Kopolovic J.

The clinical role of the PEA3 transcription factor in ovarian and breast carcinoma in effusions. Clin Exp Metastasis 2004;21(3):191-9

Doray B, Salomon R, Amiel J, Pelet A, Touraine R, Billaud M, Attie T, Bachy B, Munnich A, Lyonnet S.

Mutation of the RET ligand, neurturin, supports multigenic inheritance in Hirschsprung disease. Hum Mol Genet 1998;7:1449-1452.

Druckenbrod, N. R., and Epstein, M. L.

The pattern of neural crest advance in the cecum and colon.

Dev Biol 2005 287, 125-33.

Durbec, P. L., Larsson-Blomberg, L. B., Schuchardt, A., Costantini, F., and Pachnis, V.

Common origin and developmental dependence on c-ret of subsets of enteric and sympathetic neuroblasts.

Development 1996 122, 349-58.

Durbec P, Marcos-Gutierrez CV, Kilkenny C, Grigoriou M, Wartiovaara K, Suvanto P, Smith D, Ponder B, Costantini F, Saarma M, et al.

GDNF signalling through the Ret receptor tyrosine kinase.

Nature 1996 Jun 27;381(6585):789-93

Dutton K, Pauliny A, Lopes SS, Elworthy S, Carney TJ, Rauch J, Geisler R, Haffter P, Kelsh RN. Zebrafish colourless encodes sox10 and specifies non-ectomesenchymal neural crest fates.

Development. 2001 Nov;128(21):4113-25.

Edery P, Attie T, Amiel J et al.

Mutation of the endothelin-3 gene in the Waardenburg-Hirschsprung disease (Shah-Waardenburg syndrome).

Nat Genet 1996;12:442-444.

Edery P, Lyonnet S, Mulligan LM, Pelet A, Dow E, Abel L, Holder S, Nihoul-Fekete C, Ponder B, Munnich A.

Mutations of the RET proto-oncogene in Hirschsprung's disease.

Nature 1994a Jan 27;367(6461):378-80

Edery P, Pelet A, Mulligan LM, Abel L, Attie T, Dow E, Bonneau D, David A, Flintoff W, Jan D, et al. Long segment and short segment familial Hirschsprung's disease: variable clinical expression at the RET locus.

J Med Genet 1994b Aug;31(8):602-6

- Elworthy S, Pinto JP, Pettifer A, Cancela ML, Kelsh RN.
Phox2b function in the enteric nervous system is conserved in zebrafish and is sox10-dependent.
Mech Dev 2005 May;122(5):659-69
- Emison ES, McCallion AS, Kashuk CS, Bush RT, Grice E, Lin S, Portnoy ME, Cutler DJ, Green ED, Chakravarti A.
A common sex-dependent mutation in a RET enhancer underlies Hirschsprung disease risk.
Nature 2005 Apr 14;434(7035):857-63.
- ENCODE Project Consortium.
The ENCODE (ENCyclopedia Of DNA Elements) Project.
Science 2004 Oct 22;306(5696):636-40
- Enomoto H, Araki T, Jackman A, Heuckeroth RO, Snider WD, Johnson EM Jr, Milbrandt J.
GFR α 1-deficient mice have deficits in the enteric nervous system and kidneys.
Neuron 1998 Aug;21(2):317-24.
- Epstein, M. L., Mikawa, T., Brown, A. M., and McFarlin, D. R.
Mapping the origin of the avian enteric nervous system with a retroviral marker.
Dev Dyn 1994 201, 236-44.
- Fernandez RM, Boru G, Pecina A, Jones K, Lopez-Alonso M, Antinolo G, Borrego S, Eng C.
Ancestral RET haplotype associated with Hirschsprung's disease shows linkage disequilibrium breakpoint at -1249.
J Med Genet 2005 Apr;42(4):322-7.
- Fitze G, Appelt H, Konig I, Gorgens H, Stein U, Walther W, Gossen M, Schreiber M, Ziegler A, Roesner D, Schackert HK
Functional haplotypes of the RET proto-oncogene promoter are associated with Hirschsprung disease (HSCR).
Hum Mol Genet 2003 12:3207-3214
- Fitze G, Cramer J, Ziegler A, Schierz M, Schreiber M, Kuhlisch E, Roesner D, Schackert HK.
Association between c135G/A genotype and RET proto-oncogene germline mutations and phenotype of Hirschsprung's disease.
Lancet 2002 Apr 6;359(9313):1200-5.
- Fu M, Lui VC, Sham MH, Pachnis V, Tam PK.
Sonic hedgehog regulates the proliferation, differentiation, and migration of enteric neural crest cells in gut.
J Cell Biol 2004 Aug 30;166(5):673-84.
- Fukuda T, Kiuchi K, Takahashi M.
Novel mechanism of regulation of Rac activity and lamellipodia formation by RET tyrosine kinase.
J Biol Chem 2002 May 24;277(21):19114-21.

Furness, J. B.

"The Enteric Nervous System." 2005, Blackwell Publishing, Oxford.

Gabriel SB, Salomon R, Pelet A, Angrist M, Amiel J, Fornage M, Attie-Bitach T, Olson JM, Hofstra R, Buys C, Steffann J, Munnich A, Lyonnet S, Chakravarti A
Segregation at three loci explains familial and population risk in Hirschsprung disease.
Nat Genet 2002 31:89-93.

Garcia-Barcelo M, Ganster RW, Lui VC, Leon TY, So MT, Lau AM, Fu M, Sham MH, Knight J, Zannini MS, Sham PC, Tam PK.
TTF-1 and RET promoter SNPs: regulation of RET transcription in Hirschsprung's disease.
Hum Mol Genet 2005 Jan 15;14(2):191-204.

Garcia-Castro MI, Marcelle C, Bronner-Fraser M.
Ectodermal Wnt function as a neural crest inducer.
Science 2002 Aug 2;297(5582):848-51.

Garver KL, Law JC, Garver B.
Hirschsprung disease: a genetic study.
Clin Genet 1985 Dec;28(6):503-8.

Gershon, M. D., Kirchgessner, A. L. and Wade, P. R.
Functional anatomy of the enteric nervous system. In Physiology of the Gastrointestinal Tract, Vol. 1 (ed. L. R. Johnson), 1994 pp. 381-422. New York: Raven Press.

Goldstein AM, Brewer KC, Doyle AM, Nagy N, Roberts DJ.
BMP signaling is necessary for neural crest cell migration and ganglion formation in the enteric nervous system.
Mech Dev 2005 Jun;122(6):821-33.

de Graaff, E., Srinivas, S., Kilkenny, C., D'Agati, V., Mankoo, B. S., Costantini, F. and Pachnis, V.
Differential activities of the RET tyrosine kinase receptor isoforms during mammalian embryogenesis.
Genes Dev 2001 15,2433 -2444.

Graham A, Begbie J, McGonnell I.
Significance of the cranial neural crest.
Dev Dyn 2004 Jan;229(1):5-13

Grice EA, Rochelle ES, Green ED, Chakravarti A, McCallion AS.
Evaluation of the RET regulatory landscape reveals the biological relevance of a HSCR-implicated enhancer.
Hum Mol Genet 2005 Dec 15;14(24):3837-45

Griseri P, Bachetti T, Puppo F, Lantieri F, Ravazzolo R, Devoto M, Ceccherini I.
A common haplotype at the 5' end of the RET proto-oncogene underlies genetic susceptibility to HSCR development through reduced gene expression. Hum Mut 2005 25:189-195

Guillemot F, Joyner AL.

Dynamic expression of the murine Achaete-Scute homologue Mash-1 in the developing nervous system.

Mech Dev 1993 Aug;42(3):171-85

Hayashi, H., Ichihara, M., Iwashita, T., Murakami, H., Shimono, Y., Kawai, K., Kurokawa, K., Murakumo, Y., Imai, T., Funahashi, H., Nakao, A. and Takahashi, M.

Characterization of intracellular signals via tyrosine 1062 in RET activated by glial cell line-derived neurotrophic factor.

Oncogene 2000 19,4469 -4475

Hearn CJ, Murphy M, Newgreen D.

GDNF and ET-3 differentially modulate the numbers of avian enteric neural crest cells and enteric neurons in vitro.

Dev Biol 1998 May 1;197(1):93-105

Herbarth B, Pingault V, Bondurand N, Kuhlbrodt K, Hermans-Borgmeyer I, Puliti A, Lemort N, Goossens M, Wegner M.

Mutation of the Sry-related Sox10 gene in Dominant megacolon, a mouse model for human Hirschsprung disease.

Proc Natl Acad Sci U S A 1998 Apr 28;95(9):5161-5.

Hofstra RM, Osinga J, Tan-Sindhunata G, Wu Y, Kamsteeg EJ, Stulp RP, van Ravenswaaij-Arts C, Majoor-Krakauer D, Angrist M, Chakravarti A, Meijers C, Buys CH.

A homozygous mutation in the endothelin-3 gene associated with a combined Waardenburg type 2 and Hirschsprung phenotype (Shah-Waardenburg syndrome).

Nat Genet 1996;12:445-447.

Hofstra RM, Valdenaire O, Arch E, Osinga J, Kroes H, Loffler BM, Hamosh A, Meijers C, Buys CH. A loss-of-function mutation in the endothelin-converting enzyme 1 (ECE-1) associated with Hirschsprung disease, cardiac defects, and autonomic dysfunction.

Am J Hum Genet 1999;64:304-308.

Hofstra RM, Wu Y, Stulp RP, Elfferich P, Osinga J, Maas SM, Siderius L, Brooks AS, vd Ende JJ, Heydendaal VM, Severijnen RS, Bax KM, Meijers C, Buys CH.

RET and GDNF gene scanning in Hirschsprung patients using two dual denaturing gel systems. Hum Mutat 2000;15(5):418-29.

Holschneider AM

Hirschsprung's congenital megacolon. The concept of physiopathology and therapy

Med Welt 1982 Feb 12;33(6):210-3

Hosoda K, Hammer RE, Richardson JA, Baynash AG, Cheung JC, Giaid A, Yanagisawa M.

Targeted and natural (piebald-lethal) mutations of endothelin-B receptor gene produce megacolon associated with spotted coat color in mice.

Cell 1994 Dec 30;79(7):1267-76.

Kapur RP

Colonization of the murine hindgut by sacral crest-derived neural precursors: experimental support for an evolutionarily conserved model.

Dev Biol 2000 Nov 1;227(1):146-55

Kapur RP, Yost C, Palmiter RD.

A transgenic model for studying development of the enteric nervous system in normal and aganglionic mice.

Development 1992 Sep;116(1):167-75

Kim J, Lo L, Dormand E, Anderson DJ.

SOX10 maintains multipotency and inhibits neuronal differentiation of neural crest stem cells.

Neuron 2003 Apr 10;38(1):17-31

Kruger GM, Mosher JT, Tsai YH, Yeager KJ, Iwashita T, Gariepy CE, Morrison SJ.

Temporally distinct requirements for endothelin receptor B in the generation and migration of gut neural crest stem cells.

Neuron 2003 Dec 4;40(5):917-29.

Kusafuka T, Wang Y, Puri P.

Novel mutations of the endothelin-B receptor gene in isolated patients with Hirschsprung's disease.

Hum Mol Genet 1996 Mar;5(3):347-9.

LaBonne C, Bronner-Fraser M.

Induction and patterning of the neural crest, a stem cell-like precursor population.

J Neurobiol 1998 Aug;36(2):175-89

Lang D, Epstein JA.

Sox10 and Pax3 physically interact to mediate activation of a conserved c-RET enhancer.

Hum Mol Genet 2003 Apr 15;12(8):937-45

Le Douarin NM, Teillet MA.

The migration of neural crest cells to the wall of the digestive tract in avian embryo.

J Embryol Exp Morphol 1973 Aug;30(1):31-48

Le Douarin, N. M., and Teillet, M. A.

Experimental analysis of the migration and differentiation of neuroblasts of the autonomic nervous system and of neurectodermal mesenchymal derivatives, using a biological cell marking technique.

Dev Biol 1974 41, 162-84.

Le Douarin, N. M., and Kalcheim, C.

"The Neural Crest." 1999, Cambridge University Press, Cambridge.

- McCallion AS, Stames E, Conlon RA, Chakravarti A.
Phenotype variation in two-locus mouse models of Hirschsprung disease: tissue-specific interaction between Ret and Ednrb.
Proc Natl Acad Sci U S A 2003 Feb 18;100(4):1826-31
- Mollaaghababa RM, Hou L, Baxter LL, Loftus SK, Southard-Smith EM, Incao A, Cheng J, Pavan WJ.
Genetic evidence does not support direct regulation of EDNRB by SOX10 in migratory neural crest and the melanocyte lineage.
Mech Dev 2006 Jan 10
- Mollaaghababa R, Pavan WJ.
The importance of having your SOX on: role of SOX10 in the development of neural crest-derived melanocytes and glia.
Oncogene 2003 May 19;22(20):3024-34
- Moore, M. W., Klein, R. D., Farinas, I., Sauer, H., Armanini, M., Phillips, H., Reichardt, L. F., Ryan, A. M., Carver-Moore, K. and Rosenthal, A.
Renal and neuronal abnormalities in mice lacking GDNF.
Nature 1996 382, 76-79
- Natarajan D, Marcos-Gutierrez C, Pachnis V, de Graaff E.
Requirement of signalling by receptor tyrosine kinase RET for the directed migration of enteric nervous system progenitor cells during mammalian embryogenesis.
Development 2002 Nov;129(22):5151-60.
- Nagaya M, Kato J, Niimi N, Tanaka S, Wakamatsu N.
Clinical features of a form of Hirschsprung's disease caused by a novel genetic abnormality.
J Pediatr Surg 2002 Aug;37(8):1117-22
- Newgreen, D., Young, H. M., and Burns, A. J.
The Development of the enteric nervous system in relation to Hirschsprung's disease. In "Embryos, Genes and Birth Defects" (P. Ferretti, A. J. Copp, C. Tickle, and G. Moore, Eds.), pp. 263-300. 2006, John Wiley and Sons, Chichester.
- Oikawa T.
ETS transcription factors: possible targets for cancer therapy.
Cancer Sci 2004 95:626-633
- Paratore C, Eichenberger C, Suter U, Sommer L. Sox10 haploinsufficiency affects maintenance of progenitor cells in a mouse model of Hirschsprung disease.
Hum Mol Genet 2002;11:3075-3085.
- Pattyn A, Morin X, Cremer H, Goridis C, Brunet JF.
The homeobox gene Phox2b is essential for the development of autonomic neural crest derivatives.
Nature 1999 May 27;399(6734):366-70.

Passarge E

The genetics of Hirschsprung's disease. Evidence for heterogeneous etiology and a study of sixty-three families.

N Engl J Med 1967 Jan 19;276(3):138-43.

Pelet A, de Pontual L, Clement-Ziza M, Salomon R, Mugnier C, Matsuda F, Lathrop M, Munnich A, Feingold J, Lyonnet S, Abel L, Amiel J.

Homozygosity for a frequent and weakly penetrant predisposing allele at the RET locus in sporadic Hirschsprung disease.

J Med Genet 2005 Mar;42(3):e18

Peters-van der Sanden, M. J., Kirby, M. L., Gittenberger-de Groot, A., Tibboel, D., Mulder, M. P., and Meijers, C.

Ablation of various regions within the avian vagal neural crest has differential effects on ganglion formation in the fore-, mid- and hindgut.

Dev Dyn 1993 196, 183-94.

Pingault V, Bondurand N, Kuhlbrodt K et al. SOX10 mutations in patients with Waardenburg-Hirschsprung disease.

Nat Genet 1998;18:171-173.

Pomeranz, H. D., Rothman, T. P., and Gershon, M. D.

Colonization of the post-umbilical bowel by cells derived from the sacral neural crest: direct tracing of cell migration using an intercalating probe and a replication-deficient retrovirus. Development 1991 111, 647-55.

de Pontual L, Pelet A, Trochet D, Jaubert F, Espinosa-Parrilla Y, Munnich A, Brunet JF, Goridis C, Feingold J, Lyonnet S, Amiel J.

Mutations of the ret gene in isolated and syndromic Hirschsprung disease in human disclose major and modifier alleles at a single locus.

J Med Genet 2006 Jan 27

Potterf SB, Mollaaghababa R, Hou L, Southard-Smith EM, Hornyak TJ, Arnheiter H, Pavan WJ. Analysis of SOX10 function in neural crest-derived melanocyte development: SOX10-dependent transcriptional control of dopachrome tautomerase.

Dev Biol 2001 Sep 15;237(2):245-57

Puffenberger E, Hosoda K, Washington S, Nakao K, de Wit D, Yanagisawa M, Chakravart A. A missense mutation of the endothelin-B receptor gene in multigenic Hirschsprung's disease. Cell 1994;79:1257-1266.

Romeo G, Ronchetto P, Luo Y, Barone V, Seri M, Ceccherini I, Pasini B, Bocciardi R, Lerone M, Kaariainen H, et al. Point mutations affecting the tyrosine kinase domain of the RET proto-oncogene in Hirschsprung's disease.

Nature 1994 Jan 27;367(6461):377-8.

- Sancandi M, Griseri P, Pesce B, Patrone G, Puppo F, Lerone M, Martucciello G, Romeo G, Ravazzolo R, Devoto M, Ceccherini I
Single nucleotide polymorphic alleles in the 5' region of the RET proto-oncogene define a risk haplotype in HSCR.
J Med Genet 2003 40:714-718
- Salomon R, Attie T, Pelet A et al. Germline mutations of the RET ligand GDNF are not sufficient to cause Hirschsprung disease.
Nat Genet 1996;14:345-7.
- Schuchardt, A., D'Agati, V., Larsson-Blomberg, L., Costantini, F. and Pachnis, V. Defects in the kidney and enteric nervous system of mice lacking the tyrosine kinase receptor Ret.
Nature 1994 367,380 -383
- Serbedzija, G. N., Burgan, S., Fraser, S. E., and Bronner-Fraser, M.
Vital dye labelling demonstrates a sacral neural crest contribution to the enteric nervous system of chick and mouse embryos.
Development 1991 111, 857-66.
- Seri M, Yin L, Barone V, Bolino A, Celli I, Bocciardi R, Pasini B, Ceccherini I, Lerone M, Kristoffersson U, Larsson LT, Casasa JM, Cass DT, Abramowicz MJ, Vanderwinden JM, Kravcenkiene I, Baric I, Silengo M, Martucciello G, Romeo G.
Frequency of RET mutations in long- and short-segment Hirschsprung disease.
Hum Mutat 1997;9(3):243-9.
- Shepherd, I. T., Pietsch, J., Elworthy, S., Kelsh, R. N., and Raible, D. W.
Roles for GFRalpha1 receptors in zebrafish enteric nervous system development.
Development (2004) 131, 241-9.
- Shin MK, Levorse JM, Ingram RS, Tilghman SM.
The temporal requirement for endothelin receptor-B signalling during neural crest development.
Nature 1999 Dec 2;402(6761):496-501
- Shoba T, Dheen ST, Tay SS.
Retinoic acid influences the expression of the neuronal regulatory genes Mash-1 and c-ret in the developing rat heart.
Neurosci Lett 2002 Feb 1;318(3):129-32.
- Shoba T, Dheen ST, Tay SS.
Retinoic acid influences Phox2 expression of cardiac ganglionic cells in the developing rat heart.
Neurosci Lett 2002 Mar 15;321(1-2):41-4
- Sonnenberg-Riethmacher E, Miehle M, Stolt CC, Goerich DE, Wegner M, Riethmacher D.
Development and degeneration of dorsal root ganglia in the absence of the HMG-domain transcription factor Sox10.
Mech Dev 2001 Dec;109(2):253-65.

Southard-Smith EM, Kos L, Pavan WJ.
Sox10 mutation disrupts neural crest development in Dom Hirschsprung mouse model.
Nat Genet 1998 Jan;18(1):60-4

Stoller JZ, Epstein JA.
Cardiac neural crest.
Semin Cell Dev Biol 2005 Dec;16(6):704-15

Stolow MA, Shi YB.
Xenopus sonic hedgehog as a potential morphogen during embryogenesis and thyroid hormone-dependent metamorphosis.
Nucleic Acids Res 1995 Jul 11;23(13):2555-62

Taraviras S, Marcos-Gutierrez CV, Durbec P, Jani H, Grigoriou M, Sukumaran M, Wang LC, Hynes M, Raisman G, Pachnis V.
Signalling by the RET receptor tyrosine kinase and its role in the development of the mammalian enteric nervous system.
Development 1999 Jun;126(12):2785-97

Taraviras S, Pachnis V.
Development of the mammalian enteric nervous system.
Curr Opin Genet Dev 1999 Jun;9(3):321-7

Torfs CP, Christianson RE.
Anomalies in Down syndrome individuals in a large population-based registry.
Am J Med Genet 1998 Jun 5;77(5):431-8.

Trochet D, Bourdeaut F, Janoueix-Lerosey I, Deville A, de Pontual L, Schleiermacher G, Coze C, Philip N, Frebourg T, Munnich A, Lyonnet S, Delattre O, Amiel J.
Germline mutations of the paired-like homeobox 2B (PHOX2B) gene in neuroblastoma.
Am J Hum Genet 2004 Apr;74(4):761-4

Trochet D, O'Brien LM, Gozal D, Trang H, Nordenskjold A, Laudier B, Svensson PJ, Uhrig S, Cole T, Niemann S, Munnich A, Gaultier C, Lyonnet S, Amiel J.
PHOX2B genotype allows for prediction of tumor risk in congenital central hypoventilation syndrome.
Am J Hum Genet 2005 Mar;76(3):421-6

Verheij J and Hofstra RM: 'Shah-Waardenburg' in "Molecular basis of inborn errors of development" edited by CJ Epstein, RP Erickson and A Wynshaw-Boris. 2004 Pp. 440-446

Villanueva S, Glavic A, Ruiz P, Mayor R.
Posteriorization by FGF, Wnt, and retinoic acid is required for neural crest induction.
Dev Biol 2002 Jan 15;241(2):289-301

Wakamatsu N, Yamada Y, Yamada K, Ono T, Nomura N, Taniguchi H, Kitoh H, Mutoh N, Yamanaka T, Mushiake K, Kato K, Sonta S, Nagaya M.

Mutations in SIP1, encoding Smad interacting protein-1, cause a form of Hirschsprung disease.
Nat Genet 2001;27:369-70.

Wallace AS, Burns AJ.
Development of the enteric nervous system, smooth muscle and interstitial cells of Cajal in the human gastrointestinal tract.
Cell Tissue Res 2005 Mar;319(3):367-82

Wu JJ, Chen JX, Rothman TP, Gershon MD.
Inhibition of in vitro enteric neuronal development by endothelin-3: mediation by endothelin B receptors.
Development 1999 Mar;126(6):1161-73.

Yntema CL, Hammond WS.
The origin of intrinsic ganglia of trunk viscera from vagal neural crest in the chick embryo.
J Comp Neurol 1954 Oct;101(2):515-41

Young HM, Hearn CJ, Ciampoli D, Southwell BR, Brunet JF, Newgreen DF.
A single rostrocaudal colonization of the rodent intestine by enteric neuron precursors is revealed by the expression of Phox2b, Ret, and p75 and by explants grown under the kidney capsule or in organ culture.
Dev Biol 1998 Oct 1;202(1):67-84.

Young HM, Newgreen D.
Enteric neural crest-derived cells: origin, identification, migration, and differentiation.
Anat Rec 2001 Jan 1;262(1):1-15.

Zhu L, Lee HO, Jordan CS, Cantrell VA, Southard-Smith EM, Shin MK.
Spatiotemporal regulation of endothelin receptor-B by SOX10 in neural crest-derived enteric neuron precursors.
Nat Genet 2004 Jul;36(7):732-7

